

ELECTROWEAK PROPERTIES OF THE NUCLEON IN A CHIRAL CONSTITUENT QUARK MODEL

S. BOFFI AND M. RADICI

*Dipartimento di Fisica Nucleare e Teorica, Università di Pavia, and
INFN, Sezione di Pavia, I-27100 Pavia, Italy*

L. GLOZMAN, W. PLESSAS AND R.F. WAGENBRUNN

Institut für Theoretische Physik, Universität Graz, A-8010 Graz, Austria

W. KLINK

*Department of Physics and Astronomy, University of Iowa, Iowa City, IA 52242,
USA*

Results for all elastic electroweak nucleon form factors are presented for the chiral constituent quark model based on Goldstone-boson-exchange dynamics. The calculations are performed in a covariant framework using the point-form approach to relativistic quantum mechanics. The direct predictions of the model yield a remarkably consistent picture of the electroweak nucleon structure.

We present results for the elastic electroweak nucleon observables as a progress report of a more comprehensive programme aiming at a consistent description of the electroweak properties of baryons at low energy. The theoretical context is represented by the chiral Constituent Quark Model (CQM) based on the Goldstone-boson exchange (GBE) quark-quark interaction, that is induced by the spontaneous chiral symmetry breaking in QCD and that accurately reproduces the baryon spectrum of light and strange flavors¹. The dynamics of quarks inside the nucleon is essentially relativistic. Therefore, we have adopted the point-form realization of relativistic quantum mechanics, where the boost generators are interaction-free and make the theory manifestly covariant². The electromagnetic photon-quark interaction is assumed point-like, but in point-form the momentum delivered to the nucleon is different from the one delivered to the struck quark; hence, we will name this approach the Point-form Spectator Approximation (PFSA)³. The quark wave functions deduced from fitting the baryon spectrum are used as input and no further parameter is introduced, since quarks are considered point-like and the point-form allows for an exact calculation of all boosts required by a covariant description. Results have recently been published for electromagnetic⁴, axial⁵ and pseudoscalar³ nucleon form factors. They are summarized here in Figs. 1-2, and in Tab. 1.

The agreement with experimental data is remarkable and it indicates that




Figure 1. Proton electric and magnetic form factors. Top and middle panels: ratios of electric (G_E^p) and magnetic (G_M^p) proton form factors to the standard dipole parametrization G_D . Bottom panel: ratio of G_E^p to G_M^p . PFSA predictions of the GBE CQM (solid lines) are compared with nonrelativistic results (dashed lines) and experiment. In the top and middle panels the experimental data are from Ref. ⁶. In the bottom panel recent data from TJNAF ⁷ (filled triangles) are shown together with various older data points (see Ref. ⁷ and references therein). All the ratios are normalized to 1 at $Q^2 = 0$.

Table 1. Proton and neutron charge radii and magnetic moments as well as nucleon axial radius and axial charge. Predictions of the GBE CQM in PFSA (third column), in nonrelativistic approximation (NRIA, fourth column), and with the confinement interaction only (last column).

	Exp.	PFSA	NRIA	Conf.
r_p^2 [fm ²]	0.780(25) ⁸	0.81	0.10	0.37
r_n^2 [fm ²]	-0.113(7) ⁹	-0.13	-0.01	-0.01
μ_p [n.m.]	2.792847337(29) ¹⁰	2.7	2.74	1.84
μ_n [n.m.]	-1.91304270(5) ¹⁰	-1.7	-1.82	-1.20
$\langle r_A^2 \rangle^{\frac{1}{2}}$ [fm]	0.635(23) ¹¹	0.53	0.36	0.43
g_A	1.255 ± 0.006 ¹⁰	1.15	1.65	1.29




Figure 2. Left panel: neutron electric and magnetic form factors; in the top panel, G_E^n ; in the bottom panel, ratio of G_M^n to the standard dipole parametrization G_D , normalized to 1 at $Q^2 = 0$; solid and dashed lines as in Fig. 1; the dot-dashed line represents the PFSA results for the case with confinement only; experimental data are from Ref. ¹² (top) and Ref. ¹³ (bottom). Right panel: nucleon axial and induced pseudoscalar form factors G_A and G_P , respectively; the PFSA predictions of the GBE CQM are always represented by solid lines; in the top panel, a comparison is given to the nonrelativistic results (dashed) and to the case with a relativistic current operator but no boosts included (dot-dashed); experimental data are shown assuming a dipole parameterization with the axial mass value M_A deduced from pion electroproduction (world average: squares, Mainz experiment ¹¹: circles) and from neutrino scattering ¹⁴ (triangles); in the bottom panel, the dashed line refers to the calculation of G_P without any pion-pole contribution; the experimental data are from Ref. ¹⁵.

by a proper choice of low-energy degrees of freedom a quark model is capable of describing the spectroscopy and the low-energy dynamics of baryons at the same time. However, a more detailed comparison with data shows that there

is still room for quantitative improvements, e.g. by considering two-body electromagnetic current operators and constituent quark sizes.

References

1. L.Ya. Glozman *et al.*, *Phys. Rev. D* **58**, 094030 (1998).
2. W.H. Klink, *Phys. Rev. C* **58**, 3587 (1998).
3. S. Boffi *et al.*, [hep-ph/0108271](#), *Eur. Phys. J. A*, in press.
4. R.F. Wagenbrunn *et al.*, *Phys. Lett. B* **511**, 33 (2001).
5. L.Ya. Glozman *et al.*, *Phys. Lett. B* **516**, 183 (2001).
6. L. Andivahis *et al.*, *Phys. Rev. D* **50**, 5491 (1994); R.C. Walker *et al.*, *Phys. Rev. D* **49**, 5671 (1994); A.F. Sill *et al.*, *Phys. Rev. D* **48**, 29 (1993); G. Höhler *et al.*, *Nucl. Phys. B* **114**, 505 (1976); W. Bartel *et al.*, *Nucl. Phys. B* **58**, 429 (1973).
7. M.K. Jones *et al.*, *Phys. Rev. Lett.* **84**, 1398 (2000).
8. K. Melnikov and T. van Ritbergen, *Phys. Rev. Lett.* **84**, 1673 (2000).
9. S. Kopecky *et al.*, *Phys. Rev. Lett.* **74**, 2427 (1995).
10. D.E. Groom *et al.*, *Eur. Phys. J. C* **15**, 1 (2000).
11. A. Liesenfeld *et al.*, *Phys. Lett. B* **468**, 20 (1999).
12. T. Eden *et al.*, *Phys. Rev. C* **50**, R1749 (1994); M. Meyerhoff *et al.*, *Phys. Lett. B* **327**, 201 (1994); A. Lung *et al.*, *Phys. Rev. Lett.* **70**, 718 (1993); C. Herberg *et al.*, *Eur. Phys. J. A* **5**, 131 (1999); I. Passchier *et al.*, *Phys. Rev. Lett.* **82**, 4988 (1999); D. Rohe *et al.*, *Phys. Rev. Lett.* **83**, 4257 (1999); M. Ostrick *et al.*, *Phys. Rev. Lett.* **83**, 276 (1999); J. Becker *et al.*, *Eur. Phys. J. A* **6**, 329 (1999).
13. P. Markowitz *et al.*, *Phys. Rev. C* **48**, R5 (1993); S. Rock *et al.*, *Phys. Rev. Lett.* **49**, 1139 (1982); E.E.W. Bruins *et al.*, *Phys. Rev. Lett.* **75**, 21 (1995); H. Gao *et al.*, *Phys. Rev. C* **50**, R546 (1994); A. Lung *et al.*, *Phys. Rev. Lett.* **70**, 718 (1993); H. Anklin *et al.*, *Phys. Lett. B* **336**, 313 (1994); H. Anklin *et al.*, *Phys. Lett. B* **428**, 248 (1998).
14. T. Kitagaki *et al.*, *Phys. Rev. D* **28**, 436 (1983).
15. G. Bardin *et al.*, *Phys. Lett. B* **104**, 320 (1981); Seonho Choi *et al.*, *Phys. Rev. Lett.* **71**, 3927 (1993).